

Fecal Incident Response

(Formed stool or vomit in pool)

- ❖ Close pool to all patrons
- ❖ Remove poop or vomit with net or scoop. Sanitize net with fresh 20 ppm solution of chlorine
- ❖ Test pool for chlorine, pH, and water temperature
- ❖ Maintain pH at 7.5 or less
- ❖ Maintain water temperature at 77 F or higher
- ❖ If you use stabilized chlorine, maintain chlorine at 4 ppm for at least 25 minutes before opening pool
- ❖ If you do NOT use stabilized chlorine, maintain chlorine a 2 ppm for at least 25 minutes before opening pool
- ❖ Record incident in logbook

Fecal Incident Response

(Diarrhea in the pool)

- ❖ Close pool to all patrons
- ❖ Remove fecal material with net or scoop. Sanitize net with fresh 20 ppm solution of chlorine
- ❖ Test pool for chlorine, pH, and water temperature
- ❖ Maintain pH at 7.5 or less
- ❖ Maintain water temperature at 77 F or higher
- ❖ If you use stabilized chlorine, lower pH to 6.5 and maintain chlorine at 40 ppm for at least 30 hours before opening pool (don't use stabilized chlorine to raise chlorine levels)
- ❖ If you do NOT use stabilized chlorine, maintain chlorine a 20 ppm for at least 12 hours and 45 minutes before opening pool
- ❖ Spas shall be drained and surfaces sanitized with chlorine solution of at least 20 ppm
- ❖ Record incident in logbook